

Severe Complications of Herbal Medicines

Bobbak Vahid

Abstract

Herbal medicines are being increasingly used for treatment of variety of disorders. Herbal medicines are generally thought to lack severe side effects. Despite of the general belief, herbal medicines are known to cause serious side effects and toxicities. On the other hand, physicians' knowledge of herbal

medicines and their potential toxicities are generally limited. Neurotoxicity, cardiac toxicity, pulmonary toxicity, hepatotoxicity, and nephrotoxicity are potential severe complications of herbal medicines. The subject of severe herbal medicine complications is reviewed.

Key words: Herbal medicine; alternative medicine; side effect

Introduction

Herbal medicines are commonly used for treatment of variety of health conditions. Herbal medicines are generally believed to lack serious side effects. Lower costs compared to conventional medications are other likely attraction to these treatments. Despite of the general belief, herbal medicines can cause serious complications and even death. These complications are potentially caused by use of inherent toxic herbs, misidentification of herbs, or contamination or adulteration of herbal medicines with heavy metals or conventional medicines [1-3].

Physicians have limited knowledge regarding potential herbal medicine side effects. Physicians have little training in herbal medicines and their potential serious complications and they usually do not treat herbs in the same manner as conventional medicines. Familiarity of physicians with potential severe complications of herbal medicines is important to identify and treat these conditions [4-6]. This review discusses reported severe complications reported after use of herbal medicines.

From Thomas Jefferson University Hospital, Department of Pulmonary and Critical Care Medicine, Philadelphia, USA (Dr. Bobbak Vahid)

Author for Correspondence:

Bobbak Vahid, MD
834 Walnut Street, Suite 650
Philadelphia, PA 19107
Tel: 215 9556591
Fax: 215 9550830
Email: bobbak.vahid@mail.tju.edu

Cardiac Complications

Severe cardiac side effects can occur in patients that are taking herbal medicines that include bradycardia, heart block, and tachyarrhythmia (**Table1**) [7-23].

Erycibe henryi Prain (Ting Kung Teng) is an herb used in Chinese medicines to treat musculoskeletal discomfort. *Erycibe henryi Prain* exhibits cholinergic activities and lacrimation, salivation, rhinorrhea, diarrhea, vomiting, and miosis develop shortly after consumption. Cardiovascular manifestations are bradycardia, ventricular tachyarrhythmia, and hypotension. The management is supportive. Gastrointestinal decontamination, and in severe cases administration of intravenous atropine, could be life saving [7,8].

Complete atrioventricular block due to consumption of *Nerium oleander* containing cardiac glycoside has been reported. *Nerium oleander* is an evergreen shrub that grows in United States, China, India, Portugal, and Mediterranean region. This plant contains oleandarin and neriine that are cardiac glycosides. Despite its potential toxicity, *Nerium oleander extract* is used in mixed herbal medicines [9-11].

Caowu (*Aconitum kusnezoffii*) and Chuanwu (*Aconitum carmichaeli*) are Oriental herbal medicines that are used to treat arthritis. These herbs contain the highly toxic diterpene alkaloid: aconitine. Aconitine is a cardiotoxin. Aconitine poisoning may manifest by

gastrointestinal upset, generalized weakness, hypotension, and ventricular arrhythmias. The management of aconitine poisoning is mainly supportive. Close cardiac monitoring is necessary to detect and treat arrhythmias [12-16].

Anticholinergic syndrome may develop after ingestion of herbal teas like Angel trumpet lily tea, Lime tea mixed with *Datura innoxia*, and Paraguay tea (*Ilex paraguariensis*). Belladonna alkaloids and atropine are alkaloids responsible for the anticholinergic syndrome. Clinical manifestations include dry mucus membranes, hyperthermia, sinus tachycardia, mydriasis, cycloplegia, decreased gastrointestinal and bladder motility, and central nervous system (CNS) symptoms. Complete heart block, circulatory collapse, respiratory failure, and coma have been described with high doses. Treatment of anticholinergic syndrome is generally supportive. The use of anticholinesterase agents like tacrine hydrochloride has been suggested in severe toxicities [17-22].

Neurological Complications

Herbal medicines can cause severe neurological symptoms including weakness, seizure, and intracranial bleeding. *Cimifuga racemosa*, *Cicuta douglasii*, *Arcostaphylos uva-ursi*, *Herba ephedrae*, *Piper mysticum*, *Pausinystalia yohimbe*, *Acontium sp.*, *Ginkgo biloba*, *Strychnos nuxvomica*, *Nerium oleander*, herbal teas are known to be associated with seizures in humans. Possible mechanisms for this association include reduced seizure threshold, interaction with anti-convulsive medications, direct neurotoxicity, strychnine poisoning, and cardiovascular collapse. *Ginkgo biloba* potentially inhibits platelet activating factor and prolongs bleeding time after its consumption. Subdural hematoma and intra-cerebral bleeding have been associated with *Ginkgo biloba* [24-29].

Pulmonary Complications

Severe pulmonary complications may result after use of herbal medicines. Anaphylactic reactions, asthma exacerbation, severe interstitial pneumonitis, non-cardiogenic pulmonary edema, acute eosinophilic pneumonia, and small airway disease are discussed briefly in this section.

Peumus boldus, *Echinacea*-containing herbal remedies, and *willow bark*-containing dietary supplements are associated with anaphylactic reactions. Respiratory

support, antihistamines, epinephrine, and corticosteroids are mainstay of therapy [30-32].

Green tea has been shown to induce asthma. Green tea contains a low molecular component epigallocatechin gallate (EGCG) that is the culprit in inducing asthma. Patients with green tea-induced asthma demonstrate dose-dependent histamine release to epigallocatechin gallate challenge. IgE-mediated mechanism has also been suggested to play a role in green tea-induced asthma [33-35].

Interstitial pneumonitis in patients taking kampo medicines has been well described. Kampo medicines are mixture of several herbal agents and are used to treat a wide variety of disorders (**Table2**). The clinical presentation of kampo-induced pneumonitis is nonspecific and include fever, cough, and shortness of breath. Pneumonitis is usually detected 2 to 3 months after continues use of kampo medicines. Elevated C-reactive protein, erythrocyte sedimentation rate, serum lactic dehydrogenase, transaminases, and leukocytosis are nonspecific laboratory findings that are frequently seen in drug-induced pneumonitis. Pulmonary function testing shows carbon monoxide diffusion capacity. Chest computed tomography can findings include diffuse ground-glass opacities with patchy consolidation. The clinical course of kampo-induced pneumonitis seems to be benign. Cessation of medication may result in complete resolution of pneumonitis in several months. In more severe cases, a course of corticosteroid therapy may be reasonable [36-48].

Noncardiogenic pulmonary edema and respiratory failure have also been reported after use of kampo medicines. Although there are no data to support the use of corticosteroids, high-dose methylprednisolone has been used empirically with some success [49-51].

Acute eosinophilic pneumonia can develop after taking Shosaiko-to and Shoseiryu-to. Patients present with fever, dyspnea, hypoxemia, peripheral and BAL eosinophilia, and diffuse or peripheral pulmonary infiltrates. Resolution of pulmonary disease and hypoxemia is expected after corticosteroid therapy [52-54].

Sauropus androgynus is an Asian shrub leaf that was a commonly ingested vegetable for weight loss in Taiwan. An endemic of obstructive pulmonary disease in

young otherwise healthy women have now been attributed to *Sauropus androgynus*. These patients presented with dyspnea, obstructive pattern on pulmonary function testing, and normal chest radiographs. The obstructive lung disease is due to presence of bronchiolitis obliterans. Bronchiolitis obliterans is a pathological diagnosis. Small airways (2-4 mm) often showed segmental, partial, or completely circumferential, necrosis of the walls, with inflammatory infiltrates. Bronchiolitis obliterans is irreversible damage and conventional modalities in treatment of COPD were ineffective. Many patients developed respiratory failure and required lung transplantation [55-60].

Hepatotoxicity

Acute hepatitis has been described in patients taking herbal medicines. Extracts of *Kavalactones* are used as a remedy for anxiety and tension. *Atractylis gummifera* is a Mediterranean thistle and *Callilepis laureola* is a remedy for stomach upset and cough. The *chaparral leaf* has been used for treatment of cancer, gastrointestinal complaints, respiratory diseases and sexually transmitted diseases. *Germander* is generally consumed in tea or capsule form for weight loss. *Ma-huang* is a traditional Chinese medicine, has been commonly used in the western civilization as a weight loss aid. *Syosaiko-to* is a kampo medicine, which has been used in the treatment of chronic liver disease. Clinical manifestations usually

develop 1 to 4 months after taking the herbal medications and include nausea, vomiting, abdominal discomfort, and jaundice. Elevate transaminases are found on initial laboratory work-up. Clinical course is usually benign and most patients improve after stopping the medicines. More severe cases may progress to hepatic failure and need intensive supportive care. Orthoptic liver transplant has been performed in patients with herb-induced fulminant hepatic failure [61-68].

Nephrotoxicity

Rhizoma Rhei extracts, Aristolochia manshuriensis, Cape aloes, and Ajuga nipponensis Maniko are known nephrotoxins. Acute renal failure, acute interstitial nephritis, metabolic acidosis, rhabdomyolysis, and tubular dysfunction have been described [69-75].

Heavy Metal Poisoning

Heavy metal contamination of herbal medicine has commonly been reported. Lead, arsenic, thallium, and uranium are common contaminants. Patients may present with a variety of clinical manifestations including gastrointestinal upset, hepatitis, polyarthritis, encephalopathy, ataxia, or change of mental status. Heavy metal poisoning should be considered in patients taking herbal medications and unexplained symptoms [76-80].

TABLE 1. CARDIAC COMPLICATIONS OF HERBAL MEDICINES.

Herbal Medicine	Clinical Manifestations
Erycibe henryi Prain (Ting Kung Teng)	Cardiac: bradycardia, ventricular tachyarrhythmia, and hypotension Other: lacrimation, salivation, rhinorrhea, diarrhea, vomiting, and miosis
Nerium oleander	Cardiac: Complete Heart Block Other: Drowsiness, Tremor, Seizure
Aconitum sp.	Cardiac: Hypotension, Ventricular arrhythmias, QT-prolongation Other: Gastrointestinal Upset, Generalized Weakness
Angel trumpet lily tea Datura innoxia Ilex paraguariensis	Cardiac: Sinus Tachycardia, Complete Heart Block, Circulatory Collapse Other: Dry Mucus Membranes, Hyperthermia, Mydriasis, Cycloplegia, Constipation, Urinary Retention, CNS Symptoms.

TABLE 2. PULMONARY TOXICITIES ASSOCIATED WITH COMPLEMENTARY MEDICINES

Herbal Medicine	Pulmonary Complication
Bakumondo-to	Interstitial Pneumonia
Bofutsusho-san	Interstitial Pneumonia
Byakkokaninjin-to	Interstitial Pneumonia
Daisaiko-to	Interstitial Pneumonia
Gosha-jinki-gan	Interstitial Pneumonia
Hangeshashin-to	Interstitial Pneumonia
Kamishoyo-san	1. Interstitial Pneumonia 2. Non-cardiogenic Pulmonary Edema
Mokuboui-to	Interstitial Pneumonia
Otsuji-to	1. Interstitial Pneumonia 2. Non-cardiogenic Pulmonary Edema
Ouren-gedoku-to	Interstitial Pneumonia
Pien Tze Huang	Interstitial Pneumonia
Rikkunshi-to	Interstitial Pneumonia
Saiboku-to	Interstitial Pneumonia
Sairei-to	1. Interstitial Pneumonia 2. Non-cardiogenic Pulmonary Edema
Shosaiko-to	1. Interstitial Pneumonia 2. Non-cardiogenic Pulmonary Edema 3. Pulmonary infiltrates with eosinophilia
Shoseiryu-to	1. Interstitial Pneumonitis 2. Pulmonary infiltrates with eosinophilia
Auropus androgynus	Bronchiolitis
Green tea	Asthma exacerbation

References:

1. Basch EM, Servoss JC, Tedrow UB (2005) Safety assurances for dietary supplements policy issues and new research paradigms. *J Herb Pharmacother* 5:3-15
2. Tomlinson B, Chan TY, Chan JC, Critchley JA, But PP (2000) Toxicity of complementary therapies: an eastern perspective. *J Clin Pharmacol* 40:451-456
3. Ko RJ (2004) A U.S. perspective on the adverse reactions from traditional Chinese medicines. *J Clin Med Assoc* 67:109-116
4. Giveon SM, Liberman N, Klang S, Kahan E (2003) A survey of primary care physicians' perceptions of their patients' use of complementary medicine. *Complement Ther Med* 11:254-260
5. Silverstein DD, Spiegel AD (2001) Are physicians aware of the risks of alternative medicine? *J Community Health* 26:159-174
6. Suchard JR, Suchard MA, Steinfeldt JL (2004) Physicians knowledge of herbal toxicities and adverse herbal-drug interactions. *Eur J Emerg Med* 11:193-197
7. Huang HH, Yen DH, Wu ML, Deng JF, Huang CI, Lee CH (2006) Acute Erycibe henryi Prain ("Ting Kung Teng") poisoning. *Clin Toxicol (Phila)* 44:71-75
8. Lin CC, Chen JC (2002) Medicinal herb Erycibe henri Prain ("Ting Kung Teng") resulting in acute cholinergic syndrome. *J Toxicol Clin Toxicol* 40:185-187
9. McVann A, Havlik I, Joubert PH, Monteagudo FS (1992) Cardiac glycoside poisoning involved in deaths from traditional medicines. *S Afr Med* 81:139-141
10. Nishioka Sde A, Resende ES (1995) Transitory complete atrioventricular block associated to ingestion of Nerium oleander. *Rev Assoc Med Bras* 41:60-62
11. Dasgupta A, Datta P (2004) Rapid detection of oleander poisoning using digoxin immunoassays: comparison of five assays. *Ther Drug Monit* 26:658-663
12. Chan TY (2002) Incidence of herb-induced aconitine poisoning in Hong Kong: impact of publicity measures to promote awareness among the herbalists and the public. *Drug Saf* 25:823-828
13. Chan TY, Tomlinson B, Critchley JA, Cockram CS (1994) Herd-induced aconitine poisoning presenting as tetraplegia. *Vet Hem Toxicol* 36:133-134
14. Chan TY, Tomlinson B, Critchley JA (1993) Aconitine poisoning following the ingestion of Chinese herbal medicines: a report of eight cases. *Aust N Z J Med* 23:268-271
15. Chan TY, Tomlinson B, Chan WW, Yeung VT, Tse LK (1993) A case of acute aconitine poisoning caused by chuanwu and caowu. *J Trop Med Hyg* 96:62-63
16. Dickens P, Tai YT, But PP, Tomlinson B, Ng HK, Yan KW (1994) Fatal accidental aconitine poisoning following ingestion of Chinese herbal medicine: a report of two cases. *Forensic Sci Int* 67:55-58
17. Francis PD, Clarke CF (1999) Angel trumpet lily poisoning in five adolescents: clinical findings and management. *J Paediatr Child Health* 35:93-95
18. Pekdemir M, Yanturali S, Akay S, Alagoz G (2004) Acute anticholinergic syndrome due to Datura innoxia Miller mixed with lime tea leaves. *Vet Hum Toxicol* 46:176-177
19. Hassell LH, MacMillan MW (1995) Acute anticholinergic syndrome following ingestion of Angel's Trumpet tea. *Hawaii Med J* 54:669-670
20. Hsu CK, Leo P, Shastry D, Meggs W, Weisman R, Hoffman RS (1995) Anticholinergic poisoning associated with herbal tea. *Arch Intern Med* 155:2245-2248
21. Coremans P, Lambrecht G, Schepens P, Vanwelden J, Verhaegen H (1994) Anticholinergic intoxication with commercially available thorn apple tea. *J Toxicol Clin Toxicol* 32:589-592
22. Horowitz RS, Feldhaus K, Dart RC, Stermitz FR, Beck JJ (1996) The clinical spectrum of Jin Bu Huan toxicity. *Arch Intern Med* 156:899-903
23. Gaibazzi N, Gelmini GP, Montresor G, Canel D, Comini T, Fracalossi C, Martinetti C, Poeta ML, Ziocchi V (2002) Long QRS tachycardia secondary to Aconitum napellus alkaloid ingestion. *Ital Heart J Suppl* 3:874-877
24. Chan TY (2002) Herbal medicine causing likely strychnine poisoning. *Hum Exp Toxicol* 21:467-468
25. Benjamin J, Muir T, Briggs, Pentland B (2001) A case of cerebral haemorrhage-can Ginkgo biloba be implicated? *Postgrad Med J* 77:112-113
26. Granger AS (2001) Ginkgo Biloba precipitating epileptic seizures. *Age Ageing* 30:523-525
27. Tyagi A, Delanty N (2003) Herbal remedies, dietary supplements, and seizures. *Epilepsia* 44:228-235
28. Rowin J, Lewis SL (1996) Spontaneous bilateral subdural hematomas associated with chronic Ginkgo biloba ingestion. *Neurology* 46:1775-1776
29. Miller LG, Freeman B (2002) Possible subdural hematoma associated with Ginkgo Biloba. *J Herb Pharmacother* 2:57-63
30. Mullins RJ, Heddle R (2002) Adverse reactions associated with echinacea: the Australian experience. *Ann Allergy Asthma Immunol* 88:42-51
31. Boullata JI, McDonnell PJ, Oliva CD (2003) Anaphylactic reaction to a dietary supplement containing willow bark. *Ann Pharmacother* 37:832-835
32. Gall H, Kohler A, Peter RU (1999) Anaphylactic shock reaction to dibutylphthalate-containing capsules. *Dermatology* 199:169-170
33. Shirai T, Sato A, Hara Y (1994) Epigallocatechin gallate. The major causative agent of green tea-induced asthma. *Chest* 106:1801-1805
34. Shirai T, Sato A, Chida K, Hayakawa H, Akiyama J, Iwata M, Taniguchi M, Reshad K, Hara Y (1997) Epigallocatechin gallate-induced histamine release in patients with green tea-induced asthma. *Ann Allergy Asthma Immunol* 79:65-69
35. Shirai T, Reshad K, Yoshitomi A, Chida K, Nakamura H, Taniguchi M (2003) Green tea-induced asthma: relationship between immunological reactivity, specific and non-specific bronchial responsiveness. *Clin Exp Allergy* 33:1252-1255
36. Nakagawa A, Yamaguchi T, Takao T, Amano H (1995) Five cases of drug-induced pneumonitis due to Sho-saiko-to or interferon-alpha or both. *Nihon Kyobu Shikkan Gakkai Zasshi* 33:1361-1366
37. Katayama H, Hamada H, Yokoyama A, Kadowaki T, Ito R, Higaki J (2004) A case of interstitial pneumonia caused by gosha-jinkigan. *Nippon Ronen Igakkai Zasshi* 41:675-678
38. Hatanaka N, Yamagishi T, Kamemura H, Nakazawa I, Hirano Y, Hosaka K, Sanno K, Takahashi K (2006) A case of hepatitis and pneumonitis caused by Bofutsusyo-san herbal medicine. *Nihon Koryuiki Gakkai Zasshi* 44:335-339
39. Hatakeyama S, Tachibana A, Morita M, Suzuki K, Okano H (1997) Five cases of pneumonitis induced by sho-saiko-to. *Nihon Kyobu Shikkan Gakkai Zasshi* 35:505-510
40. Hata Y, Uehara H (2005) A case where herbal medicine sho-seiryu-to induced interstitial pneumonitis. *Nihon Koryuiki Gakkai Zasshi* 43:23-31

41. Yamawaki I, Katsura H, Taira M, Kadoriku C, Hashimoto I, Chiyotani A, Kondo M, Tamaoki J, Nagai A, Konno K (1996) Six cases with pneumonitis related to blended Chinese traditional medicines. *Nihon Kyobu Shikkan Gakkai Zasshi* 34:1331-1336
42. Takeshita K, Saisho Y, Kitamura K, Kaburagi N, Funabiki T, Inamura T, Oyamada Y, Asano K, Yamaguchi K (2001) Pneumonitis Induced by ou-gon (scullcap). *Intern Med* 40:764-768
43. Ishizaki T, Sasaki F, Ameshima S, Shiozaki K, Takahashi H, Abe Y, Ito S, Kuriyama M, Nakai T, Kitagawa M (1996) Pneumonitis during interferon and/or herbal drug therapy in patients with chronic active hepatitis. *Eur Respir J* 9:2691-2696
44. Tomioka H, Hashimoto K, Ohnishi H, Fujiyama R, Sakurai T, Tada K, Sakamoto H, Iwasaki H (1999) An autopsy of interstitial pneumonia probably induced by Sho-saikoto. *Nihon Kokyuki Gakkai Zasshi* 37:1013-1018
45. Temaru R, Yamashita N, Matsui S, Ohta T, Kawasaki A, Kobayashi M (1994) A case of drug induced pneumonitis caused by saiboku-to. *Nihon Kyobu Shikkan Gakkai Zasshi* 32:485-490
46. Suzuki K, Kinebuchi S, Sugiyama K, Satoh H, Tango M, Moriyama H, Terada M, Ooi H, Hasegawa T, Igarashi K, Satoh M, Suzuki E, Gejyo F (2002) A case of interstitial pneumonia exacerbated by kampo-induced pneumonitis. *Nihon Kokyuki Gakkai Zasshi* 40:605-611
47. Akira M, Ishikawa H, Yamamoto S (2002) Drug-induced pneumonitis: thin-section CT findings in 60 patients. *Radiology* 224:852-860
48. Maeno T, Ubukata M, Maeno Y, Suga T, Motegi M, Takayanagi N, Takahashi M, Nagai R (1997) A case of Sairei-to-induced pneumonitis diagnosed by lymphocyte stimulation test of bronchoalveolar lavage fluid. *Nihon Kyobu Shikkan Gakkai Zasshi* 35:1347-1351
49. Shiota Y, Wilson JG, Matsumoto H, Munemasa M, Okamura M, Hiyama J, Marukawa M, Ono T, Taniyama K, Mashiba H (1996) Adult respiratory distress syndrome induced by a Chinese medicine, Kamisyoyosan. *Intern Med* 35:494-496
50. Yoshida Y (2003) A non-cardiogenic type of pulmonary edema after administration of Chinese herbal medicine (shosaikoto)-a case report. *Nihon Kokyuki Gakkai Zasshi* 41:300-303
51. Sakamoto O, Ichikado K, Kohrogi H, Suga M (2003) Clinical and CT characteristics of Chinese medicine-induced acute respiratory distress syndrome. *Respirology* 8:344-350
52. Suzuki T, Higa M, Takahashi M, Saito S, Kikuchi N, Yamamuro W (2006) A case of sho-seiryu-to-induced pneumonia with a marked increase in peripheral eosinophils. *Nihon Kokyuki Gakkai Zasshi* 44:578-582
53. Soda R, Takahashi K, Tada S, Takahashi H, Okamoto S, Katagi S, Mifume T, Kajimoto K, Kimura I (1992) A case of pulmonary infiltration with eosinophilia (PIE) syndrome induced by Saiboku-To (TJ96). Detection of ECF activity in lymphocytes stimulated with TJ96. *Nihon Kyobu Shikkan Gakkai Zasshi* 30:662-667
54. Kobashi Y, Nakajima M, Niki Y, Matsushima T (1997) A case of acute eosinophilic pneumonia due to Sho-saikoto. *Nihon Kyobu Shikkan Gakkai Zasshi* 35:1372-1377
55. Hsiue TR, Guo YL, Chen KW, Chen CW, Lee CH, Chang HY (1998) Dose-response relationship and irreversible obstructive ventilatory defect in patients with consumption of Sauropus androgynus. *Chest* 113:71-76
56. Lai RS, Chiang AA, Wu MT, Wang JS, Lai NS, Lu JY, Ger LP, Roggli V (1996) Outbreak of bronchiolitis obliterans associated with consumption of Sauropus androgynus in Taiwan. *Lancet* 348:83-85
57. Chang YL, Yao YT, Wang NS, Lee YC (1998) Segmental necrosis of small bronchi after prolonged intakes of Sauropus androgynus in Taiwan. *Am J Respir Crit Care Med* 157:594-598
58. Chang H, Wang JS, Tseng HH, Lai RS, Su JM, Chang H, Wang JS, Tseng HH, Lai RS, Su JM (1997) Histopathological study of Sauropus androgynus-associated constrictive bronchiolitis obliterans: a new cause of constrictive bronchiolitis obliterans. *Am J Surg Pathol* 21:35-42
59. Wu CL, Hsu WH, Chiang CD, Kao CH, Hung DZ, King SL, Deng JF (1997) Lung injury related to consuming Sauropus androgynus vegetable. *J Toxicol Clin Toxicol* 35:241-248
60. Wang JS, Tseng HH, Lai RS, Hsu HK, Ger LP (2000) Sauropus androgynus-constrictive obliterative bronchitis/bronchiolitis--histopathological study of pneumonectomy and biopsy specimens with emphasis on the inflammatory process and disease progression. *Histopathology* 37:402-410
61. Tyler VE (1993) The honest herbal: A sensible guide to the use of herbs and related remedies. In: Foster S (Ed) New York, Coltsfood. Pharmaceutical Products Press, pp119-121
62. Borum ML (2001) Fulminant exacerbation of autoimmune hepatitis after the use of ma huang. *Am J Gastroenterol* 96:1654-1655
63. Nadir A, Agrawal S, King PD, Marshall JB (1996) Acute hepatitis associated with the use of a Chinese herbal product, ma-huang. *Am J Gastroenterol* 91:1436-1438
64. Escher M, Desmeules J, Giostra E, Mentha G (2001) Hepatitis associated with Kava, a herbal remedy for anxiety. *BMJ* 322:139
65. Hsu LM, Huang YS, Tsay SH, Chang FY, Lee SD (2006) Acute hepatitis induced by Chinese hepatoprotective herb, xiao-chai-hu-tang. *J Chin Med Assoc* 69:86-88
66. N Webb A, Hardikar W, Cranswick NE, Somers GR (2005) Probable herbal medication induced fulminant hepatic failure. *J Paediatr Child Health* 41:530-531
67. Sheikh NM, Philen RM, Love LA (1997) Chaparral-associated hepatotoxicity. *Arch Intern Med* 157:913-919
68. Mostefa-Kara N, Pauwels A, Pines E, Biour M, Levy VG (1992) Fatal hepatitis after herbal tea. *Lancet* 340:674
69. Abt AB, Oh JY, Huntington RA, Burkhart KK (1995) Chinese herbal medicine induced acute renal failure. *Arch Intern Med* 155:211-212
70. Subat-Dezulovic M, Slavic I, Rozmanic V, Persic M, Medjimurec B, Scukanec-Spoljar M (2002) Drug-induced acute tubulointerstitial nephritis: a case with elevated urinary cadmium. *Pediatr Nephrol* 17:382-385
71. Liao SC, Chiu TF, Chen JC, Lin CC (2005) Ajuga nipponensis Makino poisoning. *Clin Toxicol (Phila)* 43:583-585
72. Kwan TH, Tong MK, Leung KT, Lai CK, Poon WT, Chan YW, Lo WH, Au TC (2006) Acute renal failure associated with prolonged intake of slimming pills containing anthraquinones. *Hong Kong Med J* 12:394-397
73. Luyckx VA, Ballantine R, Claeys M, Cuyckens F, Van den Heuvel H, Cimanga RK, Vlietinck AJ, De Broe ME, Katz IJ (2002) Herbal remedy-associated acute renal failure secondary to Cape aloes. *Am J Kidney Dis* 39:E13
74. Lee CT, Wu MS, Lu k, Hsu KT (1999) Renal tubular acidosis, hypokalemic paralysis, rhabdomyolysis, and acute renal failure--a rare presentation of Chinese herbal nephropathy. *Ren Fail* 21:227-230
75. Kazama I, Matsubara M, Michimata M, Suzuki M, Hatano R, Sato H, Ito S (2004) Adult onset Fanconi syndrome: extensive tubulointerstitial lesions and glomerulopathy in the early stage of Chinese herbs nephropathy. *Clin Exp Nephrol* 8:283-287

76. Hasegawa S, Nakayama K, Iwakiri K, An E, Gomi S, Dan K, Katsumata M, Minami M, Wakabayashi I (1997) Herbal medicine-associated lead intoxication. *Intern Med* 36:56-58
77. Ernest E (2002) Heavy metals in traditional Indian remedies. *Eur J Clin Pharmacol* 57:891-896
78. Steenkamp V, Stewart MJ, Chimuka L, Cukrowska E (2005) Uranium concentration in South African herbal remedies. *Health Phys* 89:679-683
79. Saper RB, Kales SN, Paquin J, Burns MJ, Eisenberg DM, Davis RB, Phillips RS (2004) Heavy metal content of ayurvedic herbal medicine products. *JAMA* 292:2868-2873
80. Caldas ED, Machado LL (2004) Cadmium, mercury and lead in medicinal herbs in Brazil. *Food Chem Toxicol* 42:599-603